

Quenched Light Hadron Spectroscopy: Comparing the Wilson and $O(a)$ -Improved Fermion Actions

UKQCD Collaboration

presented by Alan D. Simpson

Department of Physics, University of Edinburgh,
The King's Buildings, Mayfield Road, Edinburgh, Scotland

We have studied the light hadron spectrum and decay constants for quenched QCD at $\beta = 6.2$ on a $24^3 \times 48$ lattice. We compare the results obtained using a nearest-neighbour $O(a)$ -improved (“clover”) fermion action with those obtained using the standard Wilson fermion action on the same gauge configurations. For pseudoscalar meson masses in the range 330-800 MeV, we find no significant difference between the results for the two actions. The scales obtained from the string tension and mesonic sector are consistent, but higher than those derived from baryon masses. The ratio of the pseudoscalar decay constant to the vector meson mass is roughly independent of quark mass as observed experimentally.

1. INTRODUCTION

1.1. $O(a)$ -Improved Fermion Action

Amongst the many systematic errors involved in the numerical simulation of QCD are those due to the non-zero lattice spacing, a . The standard Wilson pure gauge action differs from the continuum action by terms of $O(a^2)$, while the Wilson fermion action introduces a term of $O(a)$ in order to avoid the fermion doubling problem. Sheikholeslami and Wohlert [1] proposed a form of the fermion action which was shown by Heatlie *et al.* [2] to remove terms of $O(a)$ from matrix elements in perturbation theory. The improvement procedure described in [2] involves rotating the fermion fields and adding a local “clover” [3] term to the Wilson action, *i.e.*,

$$S_F^C = S_F^W - i \frac{\kappa}{2} \sum_{x, \mu, \nu} \bar{q}(x) F_{\mu\nu}(x) \sigma_{\mu\nu} q(x), \quad (1)$$

where $F_{\mu\nu}$ is a lattice definition of the field strength tensor.

In this paper, we present results obtained for the light hadron spectrum on a $24^3 \times 48$ lattice at $\beta = 6.2$ using both the standard Wilson action and this $O(a)$ -improved clover action.

1.2. Computational Details

The simulations were performed on the Meiko i860 Computing Surface sited in Edinburgh. This consists of 64 nodes and has a sustained performance for QCD simulations of 1 to 1.5 Gigafllops. The gauge configurations were generated using Hybrid Overrelaxation [4]. The overrelaxation was performed on a sequence of 3 $SU(2)$ subgroups of $SU(3)$. As this is a micro-canonical update, we follow every 5 overrelaxed sweeps with a Cabibbo-Marinari heatbath [5] sweep to preserve ergodicity. We allow 16800 sweeps for equilibration and separate configurations by 2400 sweeps.

The comparison of the results for the two fermion actions is performed on the same 18 configurations. For each action, we have propagators at 5 different values of the hopping parameter, κ , chosen to give roughly similar pseudoscalar masses in the range $330 \rightarrow 800$ MeV. As reported in [3], we found that the correlators obtained using the clover action were somewhat noisier and so we have smeared the clover propagators at the sink to improve the signal. We also present results for the clover action on 36 configurations with 3 κ values and a local sink.

The most computationally expensive part of this study was calculating the quark propagators. We find that red-black preconditioning improves

the efficiency by a factor of 3 and that, for the range of pseudoscalar masses investigated, Minimal Residual is typically twice as efficient as Conjugate Gradient. In agreement with the study reported by Hockney in [6], we see only a small gain from over-relaxation for MR at $\beta = 6.2$.

2. RESULTS

2.1. Fitting Procedure

Limited statistics and the amount of supercomputer time used imply that it is essential to perform a careful analysis of the data. All our propagator fits take into account correlations between timeslices. We require that the fitting region, $[t_{\min}, t_{\max}]$, satisfies the condition that small perturbations of $t_{\min}$ and $t_{\max}$ give the same mass within errors. We find that the intervals [12, 16] for local sinks and [9, 13] for smeared sinks are satisfactory. The correlated χ^2/dof varies between 0.3 and 4, which indicates both that we are taking correlations into account and that we are getting reasonable fits. Unlike the QCD-PAX collaboration [7], we find that increasing $t_{\min}$ does not significantly change the masses obtained. However, we are only able to increase $t_{\min}$ slightly before our signal becomes noisy.

The errors are estimated using a bootstrap procedure. N_{cfgs} configurations are chosen from the ensemble of N_{cfgs} configurations, allowing repetition. The same analysis is performed on this bootstrap sample as on the original N_{cfgs} configurations. We repeat this typically 1000 times and determine the errors such that the middle 68% of our bootstrap samples lie within the error bars.

2.2. Hadron Masses

Throughout the range of quark mass used, we find that m_{PS}^2 is linear with $1/2\kappa$ in accordance with PCAC. From correlated fits on 18 configurations, we obtain the following values of κ_{crit} .

$$\begin{aligned}\kappa_{\text{crit}}^W &= 0.15328^{+7}_{-4} \quad (\text{Wilson}) \\ \kappa_{\text{crit}}^C &= 0.14311^{+6}_{-3} \quad (\text{clover})\end{aligned}\quad (2)$$

The Edinburgh plot for the data is shown in Fig. 1. The results for the two actions are broadly consistent, showing a trend towards the physical value of m_N/m_V with decreasing m_{PS} .




Figure 1. Edinburgh plot for both actions. The physical and heavy quark points are marked by the left-hand and right-hand $\square$ respectively.

For the comparison of the two actions, we have concentrated on quantities which might be more sensitive to $O(a)$ effects, such as mass splittings. In Fig. 2, we plot the difference between the squares of the vector and pseudoscalar masses as a function of the square of the pseudoscalar mass in lattice units. Experimentally, this quantity is almost independent of the quark masses for both heavy-light and light-light mesons. We denote the range of experimental values by the horizontal lines, setting the scale using the string tension (*i.e.*, $a^{-1} = 2.73^{+5}_{-5}$ GeV [3]). In the chiral limit, the splitting for the Wilson action is comparable to the experimental values, while, at heavier quark mass, it is considerably smaller. For the clover action, there is a much weaker dependence on the quark mass and the values are roughly consistent with experiment.

The differences between the Δ and nucleon masses are shown in Fig. 3. The leftmost points are obtained from the chiral extrapolations of the individual masses. These values, as well as those from lighter quark masses are in good agreement with the experimental value shown.




Figure 2. $m_V^2 - m_{PS}^2$. The experimental range is indicated by the horizontal lines.

In Table 1, we present various determinations of the lattice scale. The values obtained from hadron masses are from correlated linear extrapolations to the chiral limit. The scales obtained from m_ρ agree well with the scale from the string tension, whereas those from the baryon masses, while self-consistent, are lower.

Table 1
Scales determined from various quantities.

Action	a^{-1} GeV		
	Wilson	clover	
N_{cfgs}	18	18	36
m_ρ	2.77^{+9}_{-23}	2.57^{+16}_{-16}	2.68^{+11}_{-12}
m_N	2.39^{+10}_{-12}	2.45^{+18}_{-11}	2.38^{+10}_{-15}
m_Δ	2.48^{+8}_{-15}	2.33^{+10}_{-10}	2.36^{+14}_{-13}
string tension		2.73^{+5}_{-5}	




Figure 3. Baryon mass splitting. The experimental value is marked with a $\square$.

2.3. Decay Constants

The results presented for the decay constants below are all obtained using propagators with local sinks.

2.3.1. Pseudoscalar Decay Constant

This is defined through the matrix element of the axial current

$$f_{PS} = Z_A f_{PS}^L = \frac{Z_A \langle 0 | A_4 | PS \rangle}{m_{PS}} \quad (3)$$

and our normalisation is such that the physical value is 132 MeV. We calculate f_{PS}^L using

$$f_{PS}^L = \frac{1}{m_{PS}} \sqrt{\frac{\langle A_4 A_4 \rangle}{\langle P P \rangle} |\langle 0 | P | PS \rangle|^2} \quad (4)$$

with $A_\mu = \bar{q} \gamma_\mu \gamma_5 q$ and $P = \bar{q} \gamma_5 q$. We fix m_{PS} and $\langle 0 | P | PS \rangle$ from the pseudoscalar propagator. In perturbation theory, the lattice renormalisation factors are $Z_A^W \simeq 1 - 0.13g^2$ and $Z_A^C \simeq 1 - 0.02g^2$ [8]. Using the bare coupling gives $Z_A^W \simeq 0.87$ and $Z_A^C \simeq 0.98$, whereas, if we use the “effective coupling” proposed by Lepage and Mackenzie [9], we get $Z_A^W \simeq 0.78$ and $Z_A^C \simeq 0.97$.

We have investigated the dimensionless quantity f_{PS}/m_V , and find it varies only slowly with

quark mass, in agreement with the experimental observation that f_π/m_ρ (0.17) is approximately the same as f_K/m_{K^*} (0.18). In Table 2, we show the values obtained from linear extrapolations to the chiral limit. For both actions, the chiral extrapolations are roughly consistent with experiment. The fact that the data from the clover action has larger statistical errors is compensated by the smaller uncertainty in the lattice renormalisation factor, Z_A^W . This means that we can, in principle, obtain more accurate predictions using the clover action by increasing statistics.

Table 2
Chiral extrapolations of f_{PS}/m_V .

Action	Wilson	clover	
N_{cfgs}	18	18	36
f_{PS}^L/m_V	0.21^{+2}_{-3}	0.13^{+4}_{-3}	0.15^{+2}_{-2}
$Z_A^{\text{bare}} f_{PS}^L/m_V$	0.18^{+2}_{-3}	0.12^{+4}_{-2}	0.15^{+2}_{-2}
$Z_A^{\text{eff}} f_{PS}^L/m_V$	0.16^{+2}_{-3}	0.12^{+4}_{-2}	0.15^{+2}_{-2}

2.3.2. Vector Decay Constant

We define f_V through

$$\frac{1}{f_V} \epsilon_i = \frac{Z_V}{f_V^L} \epsilon_i = \frac{Z_V}{m_V^2} \langle 0 | V_i | V \rangle \quad (5)$$

with $V_i = \bar{q} \gamma_i q$. Table 3 shows the lattice results for both actions from 18 configurations. The values depend only weakly on quark mass for both actions. The results obtained from a linear extrapolation to the chiral limit are given in the final row. Using the bare (“effective”) coupling, these correspond to $1/f_V^W = 0.39^{+2}_{-1}$ (0.33^{+1}_{-1}) and $1/f_V^C = 0.34^{+2}_{-1}$ (0.31^{+2}_{-1}). These are similar to the experimental value, $1/f_\rho = 0.28(1)$. Using the conserved vector current would remove the uncertainty in the normalisation for the Wilson action.

Table 3
 $1/f_V$ as a function of κ .

Wilson		clover	
κ	$1/f_V^L$	κ	$1/f_V^L$
0.15100	0.40^{+1}_{-1}	0.14144	0.33^{+1}_{-2}
0.15200	0.43^{+1}_{-1}	0.14226	0.36^{+2}_{-1}
0.15230	0.44^{+1}_{-1}	0.14244	0.36^{+2}_{-1}
0.15260	0.45^{+1}_{-1}	0.14262	0.37^{+2}_{-1}
0.15290	0.47^{+2}_{-1}	0.14280	0.36^{+2}_{-1}
κ_{crit}^W	0.47^{+2}_{-1}	κ_{crit}^C	0.38^{+3}_{-1}

3. CONCLUSIONS

For both actions, mass splittings and decay constants typically agree with experiment within errors. We find broadly consistent scales from different quantities, though the baryon scales appear somewhat low.

We have shown that the clover action can be used to study light hadrons. The computational overhead is small, except that the increased statistical fluctuations mean that we require smearing or more configurations for comparable errors. We see a few hints of improvement, though, in general, the results are similar to those from the standard Wilson action. This suggests that, at $\beta = 6.2$, $O(a)$ effects may be small for light hadrons.

Acknowledgements

This research is supported by the University of Edinburgh, by Meiko Limited and by the UK Science and Engineering Research Council under grants GR/G 32779, GR/H 49191 & GR/H 01069. The author also acknowledges the personal support of SERC through the award of a Postdoctoral Fellowship.

REFERENCES

1. B. Sheikholeslami & R. Wohlert, Nucl. Phys. B259 (1985) 572.
2. G. Heatlie, C.T. Sachrajda, G. Martinelli, C. Pittori & G.C. Rossi, Nucl. Phys. B352 (1991) 266.

3. *UKQCD Collaboration*, C.R. Allton *et al.*, Phys. Lett. B284 (1992) 377.
4. *UKQCD Collaboration*, Edinburgh Preprint: 92/507.
5. N. Cabibbo & E. Marinari, Phys. Lett. B119 (1982) 387.
6. G. M. Hockney, Nucl. Phys. B (Proc. Suppl.) 17 (1990) 301.
7. *QCDPAX Collaboration*, T. Yoshié, these proceedings.
8. A. Borrelli, C. Pittori, R. Frezzotti & E. Gabrielli, CERN preprint TH.6587/92.
9. G.P. Lepage & P.B. Mackenzie, Nucl. Phys. B (Proc. Suppl.) 20 (1991) 173.